

## Retrospective evaluation of lumbar puncture analysis in the pediatric intensive care unit

Retrospective analysis of pediatric meningitis

Melis Öztürk Aykaç<sup>1</sup>, Mustafa Törehan Aslan<sup>2</sup>, Nedim Samancı<sup>3</sup>

<sup>1</sup> Department of Pediatrics

<sup>2</sup> Department of Pediatrics, Division of Neonatology

<sup>3</sup> Department of Pediatrics, Division of Pediatric Intensive Care, Faculty of Medicine, Tekirdağ Namık University, Tekirdağ, Turkey

### Abstract

**Aim:** Meningitis is a central nervous system infectious disease with inflammation of the leptomeninges and subarachnoid space. There is still a severe mortality and morbidity from meningitis. Therefore, it is essential to make an early diagnosis and start empirical treatment for the causative agent as soon as possible. **Material and Methods:** Seventy patients diagnosed with meningitis in our hospital were included in our study. The anamnesis, physical examination, laboratory results, and treatment of the patients were retrospectively reviewed.

**Results:** The most common complaint of our patients was fever, with 84.3%. Meningeal irritation finding was positive in 86% of patients aged two years and older, while this rate was 22.4% in patients under two years of age. According to the meningeal irritation findings, a statistically significant difference was found in terms of neutrophil, lymphocyte, and platelet values ( $p < 0.05$ ). Neutrophil levels of those with positive signs of meningeal irritation were significantly higher than in those without symptoms. Lymphocyte and platelet values of those with positive signs of meningeal irritation were substantially lower than in those without. The median values were 11.5 g/dL for hemoglobin, 11220/uL for leukocytes, 321500/uL for platelets, 25.9 mg/dl for CRP, 29 mm/h for ESR, 61.2 mg/dL for CSF glucose, 37.4 mg/dL for CSF protein, and 100 for cell count.

**Discussion:** In 4 of our patients, the CSF culture detected the causative microorganism. In 3 of the culture-negative patients, the causative pathogen was detected in CSF PCR. The most frequently preferred empirical antimicrobial treatment initiated for patients is the combination of vancomycin and a third-generation cephalosporin.

### Keywords

Meningitis, Fever, Meningeal Irritation Signs, Cerebrospinal Fluid

DOI: 10.4328/ACAM.21832 Received: 2023-07-26 Accepted: 2023-08-28 Published Online: 2023-12-29 Printed: 2024-03-01 Ann Clin Anal Med 2024;15(3):145-148

Corresponding Author: Nedim Samancı, Department of Pediatrics, Division of Pediatric Intensive Care, Faculty of Medicine, Tekirdağ Namık University, Tekirdağ, Turkey.

E-mail: samancitr@yahoo.com P: +90 532 295 71 98

Corresponding Author ORCID ID: <https://orcid.org/0000-0002-3947-3492>

This study was approved by the Ethics Committee of Tekirdag Namik Kemal University Non-Interventional Clinical Research Ethics Committee (Date: 2021-03-30, No: 2021.66.03.06)

## Introduction

Meningitis is a central nervous system infectious disease that describes the inflammation of the meninges and the cerebrospinal fluid in the subarachnoid space resulting in severe mortality and morbidity [1, 2]. Bacteria, viruses, and other infectious agents can cause meningitis [1]. Clinical findings and morphological, chemical, and microbiological examination of the cerebrospinal fluid can reveal the underlying cause. The main goal in treating meningitis is to diagnose and start treatment as soon as possible, to prevent or minimize the complications that may develop [3, 4]. This research was carried out in the pediatric intensive care unit of our hospital to evaluate the clinical findings, blood tests, chemical and microbiological examinations in the cerebrospinal fluid and to examine the demographic characteristics, application complaints, examination findings, and test results of the patients diagnosed with meningitis, the compatibility of the biochemical and microbiological parameters obtained from these results, and the contribution of all these results to the diagnosis [1, 5].

## Material and Methods

Demographic characteristics, anamnesis, and physical examination findings, biochemical and microbiological data of 70 patients aged one month to 18 years who underwent lumbar puncture with the diagnosis of meningitis in Tekirdağ Namik Kemal University Hospital, Department of Pediatrics, Pediatric Intensive Care Unit were included in the study by retrospectively examining. Ethical approval was obtained from Tekirdağ Namik Kemal University Non-Interventional Clinical Research Ethics Committee (Study protocol number: 2021.66.03.06, Date: 30.03.2021).

### Statistical analyses

Statistical analyses were performed using a package program called SPSS (IBM SPSS Statistics 24). Frequency tables and descriptive statistics were used to interpret the findings. Parametric methods were used for measurement values suitable for a normal distribution. Using parametric methods, the "Independent Sample-t" test (t-table value) method was used to compare the measurement values of two independent groups. Non-parametric methods were used for the measurement values that did not conform to the normal distribution. By non-parametric methods, the "Mann-Whitney U" test (Z-table value) method was used to compare the measurement values of two independent groups. The Spearman correlation coefficient was used to examine the relationships of two quantitative variables that do not have a normal distribution. "Pearson-x<sup>2</sup>" crosstabs were used to explore the relationships between two qualitative variables.

## Results

It was determined that the mean age of the patients was  $6.11 \pm 5.33$  (years), and 27 (38.6%) were in the  $\leq 1$  age group. It was determined that 40 patients (57.2%) applied in the spring and summer months; 41 (58.6%) were male, 29 (41.4%) were female, and 5 (7.1%) had ventriculoperitoneal shunts (Table 1). A statistically significant difference was found in admission complaints regarding age (years) and head and neck pain status

( $Z = -6.431$ ;  $p = 0.0001$ ). A statistically significant difference was found in age (years) according to MIB status at the first examination ( $Z = -5.829$ ;  $p = 0.0001$ ). It was observed that the age of those with MIB was significantly higher than that of those without. A statistically significant correlation was found between the patient group under two years of age and the MIB status ( $\chi^2 = 25.677$ ;  $p = 0.0001$ ). A statistically significant correlation was found between the patient group under two years of age and anterior fontanel status ( $\chi^2 = 35.758$ ;  $p = 0.0001$ ). It was determined that the anterior fontanel of 17 individuals (63%) under the age of two years was open. Moreover, it was determined that those in the  $< 2$  age group were predominantly open (bomb and pulsatile), while those in the age group over two years old were all closed. Thirty-one of the patients (44.3%) had very intense leukocytes in the CSF culture, 64 (91.4%) had PMNL as the dominant cell, 66 (94.3%) had no growth in the CSF culture, four patients (5.7%) had change in their culture, two patients (50.0%) had change in CSF culture, pneumococcal growth was found in 3 (4.2%) of meningitis. There was no statistically significant difference in terms of leukocytes, neutrophils, lymphocytes, platelets, and

**Table 1.** Distribution of findings related to patients

Variable (n=70)	n	%
Application season		
Spring	20	28,6
Summer	20	28,6
Autumn	16	22,8
Winter	14	20
Age classes [ $\bar{X} \pm S.S. \rightarrow 6.11 \pm 5.33$ (years) ]		
$\leq 1$ year	27	38,6
2-5 years	6	8,6
6-9 years	15	21,4
$\geq 10$ years	22	31,4
Gender		
Female	29	41,4
Male	41	58,6
VP shunt		
No	65	92,9
Yes	5	7,1

**Table 2.** Distribution of biochemical and microscopic examination results of the patients

Variable	Mean	S.D.	Median	Min.	Max.
Hemoglobin (g/dL)	11,6	1,48	11,5	8,2	15
Leukocytes (1/uL)	15081,42	20721,27	11220	2010	17800
Neutrophil (1/uL)	8690,57	6262,54	7110	1180	32000
Lymphocyte (1/uL)	3051,71	1984,03	2435	420	8250
Platelet (1/uL)	345014,29	131731,89	321500	117000	753000
CRP*(mg/dl)	68,34	90,93	25,9	0,1	432
ESR**(mm/hr)	36,38	27,08	29	4	121
CSF*** Glucose (mg/dl)	59,94	19,39	61,2	2	124
CSF protein (mg/dl)	77,86	91,94	37,4	13,7	553
Number of cells/mm <sup>3</sup>	910	2613,29	100	10	17000

CRP\*: C-Reactive Protein, ESR\*\*: Erythrocyte Sedimentation rate, CSF\*\*\*: Cerebrospinal Fluid

C-Reactive Protein (CRP) according to the presence of fever in complaints at admission ( $p > 0.05$ ). A statistically significant difference was found in erythrocyte sedimentation rate (ESR) values according to the presence of fever in complaints at admission ( $Z = -2.442$ ;  $p = 0.015$ ). It was observed that the ESR values in those who had an uproar in the application complaint were significantly higher than in those who did not have a fever. There was no statistically significant difference in leukocytes, neutrophils, CRP, and ESR values according to the presence of head and neck pain in the presenting complaint ( $p > 0.05$ ).

A statistically significant difference was found in lymphocyte and platelet values according to head and neck pain status ( $p < 0.05$ ). There was no statistically significant difference in the admission complaints regarding nausea, vomiting, sleepiness, leukocytes, neutrophils, platelets, CRP, and ESR values ( $p > 0.05$ ). There was no statistically significant difference in leukocytes, neutrophils, lymphocytes, CRP, and ESR according to seizure status at admission ( $p > 0.05$ ). However, a statistically significant difference was found in platelet values ( $t = -2.135$ ;  $p = 0.036$ ). Platelet values of those with seizure complaints were significantly higher than in those without seizures. At the first examination, there was no statistically significant difference in leukocytes, CRP, and ESR according to the meningeal irritation findings in the patients ( $p > 0.05$ ). According to the meningeal irritation findings, a statistically significant difference was found in terms of neutrophil, lymphocyte, and platelet values ( $p < 0.05$ ). Neutrophil levels in those with positive signs of meningeal irritation were significantly higher than in those without symptoms of meningeal irritation. A positive, weak, and statistically significant correlation was found between the patients' leukocyte values and CRP values ( $r = 0.399$ ;  $p = 0.001$ ). As CRP increases, leukocytes will increase. Likewise, it was observed that leukocyte values decreased as CRP decreased. A positive, weak, statistically significant relationship was found between ESR and CRP ( $r = 0.440$ ;  $p = 0.0001$ ). As CRP increases, ESR will increase. When the distribution of the patients' biochemical and microscopic examination results was examined, the median cell number was  $f00/mm^3$ , the median CSF protein was  $61.2 \text{ mg/dl}$ , and the median CRP was  $25.9 \text{ mg/dl}$ . The median leukocyte count was 11220, which is detailed in Table 2.

## Discussion

Childhood meningitis has been one of the most feared infectious diseases since ancient times [1]. Acute bacterial meningitis, which continues to cause death and disability in children, is a neurological emergency, especially in developing countries with weak immunization studies [6]. Despite advances in antimicrobial treatments, the rate of development of acute bacterial meningitis in some developing countries varies between 16-32% and is still very high [7, 8].

Most of our study's patients (38.6%) were one year old or younger. We think that this result is due to the immaturity of the immune response and the lumbar puncture to be performed primarily when no fever focus can be found in a patient younger than one year of age and the vaccine doses have not been completed yet. It has been stated that the only risk factor is age under 1 year, and the mortality rate under 1 year of age is

high; therefore, this age group should be closely monitored. The same study emphasized that the reasons behind the uncertain seasonal bacterial meningitis patterns should be investigated further. Considering the application dates of the 70 patients in our study, according to the seasonal distribution we made, the total number of patients admitted in the spring and summer months (57.2%) is higher than the number of patients admitted in the autumn and winter months (42.8%). However, no significant difference was observed between the distributions when the seasons were evaluated alone.

The most common finding in bacterial meningitis is fever [8]. When the symptoms of 70 patients in our study were examined, the most common symptom was fever in 59 patients (84.3%). The most common symptom after turmoil is head and neck pain in 34 (48.6%) patients. A statistically significant difference was found in age (years) according to head and neck pain status in admission complaints. Those with head and neck pain are significantly older than those without. Considering both the relationship of anterior fontanel status with age and the requirement that children must be above a certain age to define head and neck pain as a complaint, it is customary to be statistically significant.

A statistically significant difference was found in admission complaints regarding age (years) and head and neck pain status ( $Z = -6.431$ ;  $p = 0.0001$ ). It was observed that the ages of those with head and neck pain were significantly higher than those without. A statistically significant difference was found in age (years) according to MIB status at the first examination ( $Z = -5.829$ ;  $p = 0.0001$ ). It was observed that the ages of those with MIB were significantly higher than those without.

In the study by Michos et al., in which they examined 506 patients with aseptic meningitis in Greece, the median cell number in the CSF was found to be  $201/mm^3$ , and PMNL was the dominant cell in 58.3% of the patients [9]. PMNL predominates in the CSF in approximately 90% of patients with bacterial meningitis. PMNL predominates in the early period in 20-75% of cases with viral meningitis [10]. In the CSF cell analysis of 70 patients in our study, the median cell number was 100 per  $mm^3$ , and in the CSF analysis of 64 (94.3%) of the patients, PMNL was dominant when cell typing was performed. *S.pneumoniae* growth was detected in 2 of 4 patients with development.

In our study, five patients (7.1%) had ventriculoperitoneal shunts, and all meningitis patients with VP shunts were younger than one year old. Enterococci were found in CSF culture in 1 patient with VP shunt, and Methicillin-Resistant *Staphylococcus hominis* growth was detected in CSF culture in 1 patient. However, since the number of patients with VP shunt is limited, it is difficult to make a judgment.

The diagnosis of bacterial meningitis is confirmed by microbiological testing of CSF, which includes cytochemical analysis, cell count, microscopy, and culture [11, 12]. However, antibiotics before the lumbar puncture procedure may reduce the culture yield and cause diagnostic uncertainty [12, 13].

In our study, when the symptoms and signs were compared with some biochemical parameters, ESR values in patients with fever were statistically significantly higher than in those without fever. No statistically significant difference was found

between the patients with and without fever for values of other leukocytes, neutrophils, lymphocytes, and CRP. Lymphocyte and platelet values in those with head and neck pain were statistically significantly lower than in those without head and neck pain. It was observed that neutrophil values in patients with seizures were statistically significantly higher than in those without seizures. Neutrophil values in those with meningeal irritation findings were statistically significantly higher than in those without, and lymphocyte and platelet values were lower.

### Conclusion

The mean age of the patients was  $6.11 \pm 5.33$  (years), and 37.2% were one year old or younger. When patients' admission symptoms and physical examination findings are examined, the most common symptom is fever, with a rate of 84.3%. ESR values in patients with fever at admission are statistically significantly higher than in those without. No statistically significant difference was found between the patients with and without fever for other values of leukocytes, neutrophils, lymphocytes, and CRP. The most common symptom after turmoil is head and neck pain, which is seen in 48.6%, and a statistically significant difference was found in age (years) according to head and neck pain status in admission complaints ( $Z=-6.431$ ;  $p=0.0001$ ). A statistically significant difference was found in lymphocyte and platelet values according to head and neck pain status ( $p<0.05$ ). The lymphocyte and platelet values in those with head and neck pain were significantly lower than in those without head and neck pain.

The meningeal irritation findings found a statistically significant difference in neutrophil, lymphocyte, and platelet values ( $p<0.05$ ).

Growth was observed in the CSF culture of four patients, and *S. pneumoniae* growth was detected in 2 of them. In the other two patients with development, VP shunt was found. Enterococcus growth was seen in one of the CSF cultures, and Methicillin-Resistant *Staphylococcus hominis* growth was found in the CSF culture of the other patient. Five of the patients included in our study had VP shunt, and all of the patients with VP shunt were under one year old. In patients in whom the CSF PCR was studied, no growth was observed in the CSF culture of 3 patients, and *N. meningitidis* was found in one patient, Hib in one, and Parechovirus in the other in the CSF PCR results. The causative pathogen was detected by CSF culture and PCR methods in only seven of 70 patients in our study. Although the etiology cannot be determined, the patients are those whose meningitis is suspected by anamnesis, physical examination, and laboratory findings, and the diagnosis of meningitis is confirmed by the CSF's cell count and the CSF's biochemical analysis. For this reason, especially unexplained fever, headache, vomiting, and positive signs of meningeal irritation on physical examination, the presence of a bulging and pulsatile fontanelle in a child with open anterior fontanelle should bring meningitis to mind in the diagnosis, and especially if there are no contraindications, the presence of meningitis should be considered. If suspected, acyclovir treatment for HSV should be started.

### Scientific Responsibility Statement

The authors declare that they are responsible for the article's scientific content including study design, data collection, analysis and interpretation, writing, some of the main line, or all of the preparation and scientific review of the contents and

approval of the final version of the article.

### Animal and Human Rights Statement

All procedures performed in this study were in accordance with the ethical standards of the institutional and/or national research committee and with the 1964 Helsinki Declaration and its later amendments or comparable ethical standards.

**Funding:** None

### Conflict of Interest

The authors declare that there is no conflict of interest.

### References

- Hasbun R. Progress and Challenges in Bacterial Meningitis: A Review. *Jama*. 2022;328(21):2147-54.
- Van Hentenryck M, Schroeder AR, McCulloh RJ, Stave CD, Wang ME. Duration of Antibiotic Therapy for Bacterial Meningitis in Young Infants: A Systematic Review. *Pediatrics*. 2022;150(5):e2022057510 DOI: 10.1542/peds.2022-057510.
- Silva GD, Guedes BF, Junqueira IR, Gomes HR, Vidal JE. Diagnostic and therapeutic approach to chronic meningitis in Brazil: A narrative review. *Arq Neuropsiquiatr*. 2022;80(11):1167-77.
- Mijovic H, Sadarangani M. To LP or not to LP? Identifying the Etiology of Pediatric Meningitis. *Pediatr Infect Dis J*. 2019;38(6S Suppl. 1):S39-42.
- de Almeida SM, Castoldi JR, Riechi SC. Comparison between cerebrospinal fluid biomarkers for differential diagnosis of acute meningitis. *Diagnosis (Berl)*. 2023; 10(3):298-308 DOI: 10.1515/dx-2023-0013.
- Kuti BP, Bello EO, Jegede TO, Olubosede O. Epidemiological, clinical and prognostic profile of childhood acute bacterial meningitis in a resource poor setting. *J Neurosci Rural Pract*. 2015;6(4):549-57.
- Jarousha AM, Afifi AA. Epidemiology and Risk Factors Associated with Developing Bacterial Meningitis among Children in Gaza Strip. *Iran J Public Health*. 2014;43(9):1176-83.
- Nickerson JW, Attaran A, Westerberg BD, Curtis S, Overton S, Mayer P. Fatal Bacterial Meningitis Possibly Associated with Substandard Ceftriaxone--Uganda, 2013. *MMWR Morb Mortal Wkly Rep*. 2016;64(50-51):1375-7.
- Michos AG, Syriopoulou VP, Hadjichristodoulou C, Daikos GL, Lagona E, Douridas P, et al. Aseptic meningitis in children: Analysis of 506 cases. *PLoS one*. 2007;2(7):e674.
- Kanra G, Ceyhan M, Kara A. Meningitis II: Clinical findings. *Çocuk Sağlığı ve Hastalıkları Dergisi/ Journal of Child Health and Diseases*. 2003;46(2):128-38.
- Polage CR, Cohen SH. State-of-the-Art Microbiologic Testing for Community-Acquired Meningitis and Encephalitis. *J Clin Microbiol*. 2016;54(5):1197-202.
- Khumalo J, Nicol M, Hardie D, Muloiwa R, Mtshana P, Bamford C. Diagnostic accuracy of two multiplex real-time polymerase chain reaction assays for the diagnosis of meningitis in children in a resource-limited setting. *PLoS one*. 2017;12(3):e0173948.
- Abdelrahim NA, Mohammed N, Evander M, Ahlm C, Fadl-Elmula IM. Viral meningitis in Sudanese children: Differentiation, etiology and review of literature. *Medicine*. 2022;101(46):e31588.

### How to cite this article:

Melis Öztürk Aykaç, Mustafa Törehan Aslan, Nedim Samancı. Retrospective evaluation of lumbar puncture analysis in the pediatric intensive care unit. *Ann Clin Anal Med* 2024;15(3):145-148

This study was approved by the Ethics Committee of Tekirdag Namik Kemal University Non-Interventional Clinical Research Ethics Committee (Date: 2021-03-30, No: 2021.66.03.06)